

1


# KHC Office Hours Population Health Strategies for Cardiovascular and Kidney Disease Risk Reduction April 23, 2025 Elizabeth Montgomery, National Kidney Foundation Michelle Falgoust, National Kidney Foundation Tiffany Scheffler, National Kidney Foundation KANSAS MEDICAL Kansas Healthcare COLLA BORATIVE KANSAS Healthcare COLLA BORATIVE

2


### DISCLAIMER

All Kansas Healthcare Collaborative, Inc. ("KHC") content and information in KHC's publications, presentations, and on KHC's website ("KHC Content") are for informational and educational purposes only. KHC Content does <u>not</u> create any type of relationship with, or duty to, the reader, attendee, or user. KHC Content does <u>not</u> constitute legal, tax, business, professional, or personal advice. KHC disclaims any and all liabilities and warranties, express or implied, arising from users' access and use of KHC Content.


ansas Health




4




5







7



8







10




11







13




14






16




17


Population Health for CKD and Diabetes: Lessons from the Indian Health Service	Measures for CKD Care included in IHS Diabetes Care & Outcomes Audit Intervention		
"The 54% reduction in incidence occurred in this population during a 20-year period despite per capita health expenditures equaling only "40% of that spent in the US civilian population. Although one might expect such a dramatic decrease in disease in this high-risk disadvantaged population to be associated with novel and costly new therapies, the medical interventions implemented by the HIS were routine: glucose control, blood pressure control, and use of renin-angiotensin-aldosterone system (RAAS) antagonists in appropriate patients. However, a systematic population-based approach was instituted to implement this evidence-based care."	Measure	Baseline	Impact
	Improve average Hemoglobin A1C among people with DM	10%	8.1%
	Continue blood pressure control among people with DM and CKD	133/76 mmHg	133/76 mmHg
	Increase Urine Albumin-Creatinine Ratio Testing for early detection	50%	62%
	Increase use of Ace Inhibitors (ACE) and Angiotensin Receptor Blockers (ARB)	42%	73%
	Intensive focus on mitigating health-related social needs and lifestyle impacts included in this intervention		
	Narva A. Population Health for CKD and Diabetes: Lessons From&tra0, the Indian Health Service. <i>American Journal of Kidney Diseases</i> . 2018;71(3):407-411.		



19




20






22



23







25




26





28



29





31

### **Upcoming Education and Important Dates**

- 4/25 KHA Grants at a Glance
- 4/30 NARHC Office Hours
- 5/1 Noon Briefing Self-funded Plans Simplified
- 5/6 Quality Corner Call
- . 5/20 KHC CAP Lunch and Learn RHC Compliance Are you survey ready?
- 5/20 Critical Marketing KHSC Introductory Webinar
   7/15 KHC CAP Lunch and Learn RHC Billing Basics
- · KHC Summit on Quality August 7th

32





34

